

# AFTER CARE

## Moderna Spikevax® Bivalent Original/Omicron BA.4/5 COVID-19 Vaccine

Rev. July 2023

### What should I/my child do right after receiving the vaccine?

- **Wait for at least 15 minutes** after receiving the vaccine. A longer waiting time of 30 minutes may be recommended
- if there is concern about a possible vaccine allergy. Though uncommon, fainting or allergic reactions can occur after vaccination. Symptoms of an allergic reaction include hives (bumps on the skin that are often very itchy), swelling of the face, tongue or throat, or difficulty breathing. The clinic staff are prepared to manage these events should they occur.
- You/your child may be asked to wait in the clinic or outside the clinic (such as in your vehicle) if an adult is with you/them. If asked to wait outside, inform the health care provider if you do not have a warm, dry place to wait.
- **Inform a health care provider** at the clinic if you/your child feel unwell while waiting.
- If waiting inside the clinic, be sure to **leave your/your child's mask on** if at all possible and **remain at least 2 metres away** from anyone not part of your household.
- Use the **alcohol-based hand rub to clean your/your child's hands** before leaving the clinic.
- **Do not operate a vehicle** or other form of transportation for at least 15 to 30 minutes after being vaccinated (as advised by the health care provider) or if you are feeling unwell.

### What should be expected in the next few days?

- Side effects can develop within 3 days after receiving the vaccine. Although these side effects are not serious to your/your child's health, they may make you/your child feel unwell for a day or two; they will go away on their own.
  - » The most common side effect is pain where the needle was given; swelling where the needle was given may also occur. A cool, damp cloth or wrapped ice pack where the vaccine was given may help.

- » Some local reactions may occur 7 days or more after immunization and include a rash, tenderness and extensive swelling where the injection was given. This is a known reaction that resolves on its own within 7-10 days.
- » Other symptoms may include tiredness, headache, muscle pain, joint pain, chills or fever. Enlarged lymph nodes (swollen glands) that last for several days may also occur.
- If needed, pain or fever medication (such as acetaminophen or ibuprofen) may help with pain or fever. Check with your health care provider if you need advice about medication.
- Serious side effects after receiving the vaccine are rare. However, **should you/your child develop any serious symptoms or symptoms that could be an allergic reaction, call 9-1-1 right away.**
- Symptoms of an allergic reaction include:
  - » Hives (bumps on the skin that are often very itchy)
  - » Swelling of the face, tongue or throat
  - » Difficulty breathing
- Very rare cases of myocarditis (inflammation of the heart) and/or pericarditis (inflammation of the outer lining of the heart) following immunization with this vaccine have been reported in those 12+ years old. Go to Emergency or call 911 right away if you/your child have any of the following symptoms any time after vaccination:
  - » chest pain
  - » shortness of breath
  - » irregular heartbeat, a very fast or pounding heartbeat
- Very rare reports of Bell's palsy (weakness or paralysis on one side of the face that is often temporary) following immunization have been reported in a small number of people 12+ years old in Canada and internationally. Symptoms usually resolve within a month.

- If you have any concerns about these or other symptoms you develop after receiving the vaccine, call 811. If you/your child received a vaccine from a pharmacist, please consult them first when possible.
- Tell the person providing the next dose about any side effects you/your child experienced after a previous dose.

### When should I/my child return for the next vaccine dose?

- **Primary series:** Return for the second dose at the optimal time of 8 weeks after the first dose (minimum 21 days). Eligible high-risk individuals should receive 3 doses as their primary series with 4-8 weeks between each dose.
- **Booster dose for ages 6 years and older:** After the last dose in a primary series, it is recommended to return for one bivalent booster dose after 6 months. This is important to ensure the vaccine continues to work well.
- Seniors, immune compromised adults and adults living in personal care homes and long-term care facilities are recommended to get a second bivalent booster dose 6 months after their first bivalent booster dose.
- A person is considered up-to-date on their vaccinations if they have received the last dose they are eligible for.
- Ask your healthcare provider or visit <https://www.saskatchewan.ca/covid19-vaccine> for booster dose information.

### Things to remember

- **Continue to follow the recommendations of local public health officials** to prevent spread of COVID-19, which may include wearing a mask, staying at least 2 metres from others and limiting/avoiding contact with others outside your household.
- **Keep this sheet or other immunization record in a safe place.** You can also download the CANImmunize app to keep track of this and other vaccines. Sign up for MySaskHealthRecord, a secure website that enables Saskatchewan citizens to quickly and easily access their personal health information including immunization history.