

SCREENING

Novavax Nuvaxovid® COVID-19 Vaccine

Rev. March 28, 2023

1. Are you feeling ill today?

- » Do not attend a public immunization clinic if you have any new or worsening respiratory symptoms (fever, cough, sore throat, runny nose).

2. Have you had a previous COVID -19 infection?

- » Immunization with a COVID-19 vaccine dose in a primary series should be given 3 months after infection.
- » Immunization with a COVID-19 vaccine booster dose should be given 6 months after infection.
- » More time between infection and vaccination is recommended to ensure a strong immune response.

3. Are you pregnant, could you be pregnant or are you planning on becoming pregnant before receiving both doses of the vaccine?

- » Only mRNA COVID-19 vaccines should be offered during pregnancy unless there are contraindications. Other COVID-19 vaccines should only be offered if there are allergies to mRNA vaccine ingredients or mRNA vaccine is not readily available.
- » There is a pregnancy exposure registry that monitors pregnancy outcomes in women exposed to NUVAXOVID during pregnancy. Women who are vaccinated with NUVAXOVID during pregnancy are encouraged to enroll in the registry by visiting <https://c-viper.pregistry.com/>.

4. Are you breastfeeding?

- » Only mRNA COVID-19 vaccines should be offered when breastfeeding unless there are contraindications. Other COVID-19 vaccines should only be offered if there are allergies to mRNA vaccine ingredients or mRNA vaccine is not readily available.

5. Are you allergic to any components in the vaccine including polysorbate 80?

- » Polysorbate 80 is a component of this vaccine.
- » Polysorbate 80 can be found in medical preparations such as vitamin oils, tablets, anticancer agents, vaccines and cosmetics.

6. Do you have a history of myocarditis (inflammation of the heart) or pericarditis (inflammation of the lining of the heart) not related to COVID-19 immunization?

- » This is not a contraindication to being immunized with COVID-19 vaccines.

7. Do you have a history of myocarditis (inflammation of the heart) or pericarditis (inflammation of the lining of the heart) after getting a previous dose of COVID-19 vaccine?

- » As a precautionary measure, additional protein-based COVID-19 vaccine doses should be deferred in individuals who developed myocarditis or pericarditis following a dose of a COVID-19 vaccine until more information is available. If an individual is at high risk of being infected with COVID-19 due to community transmission or of severe illness due to an underlying condition, then a decision to get an additional dose should be made in consultation with the individual's physician (cardiologist if possible) with the individual's informed consent.

8. Do you have any problems with your immune system or are you taking any medications/treatment that can affect your immune system (e.g., high dose steroids, chemotherapy)?

- » Only mRNA COVID-19 vaccines should be offered to immune compromised persons unless there are contraindications. Other COVID-19 vaccines should only be offered if there are allergies to mRNA vaccine ingredients or mRNA vaccine is not readily available.
- » The vaccine immune response may not be as strong for people with immune system conditions due to disease or treatment compared to people who do not have immune system conditions.
- » Stem cell transplant recipients MUST consult their specialist prior to immunization with COVID-19 vaccine.
- » For transplant recipients, it is preferred to contact your health care provider to determine the optimal time to receive the COVID-19 vaccine.

9. Do you have an autoimmune disease?

- » Only mRNA COVID-19 vaccines should be offered to those with an autoimmune condition unless there are contraindications. Other COVID-19 vaccines should only be offered if there are allergies to mRNA vaccine ingredients or mRNA vaccine is not readily available.
- » For persons with any autoimmune condition that involves the neurological system it is preferred they discuss this with their primary physician or specialist prior to immunization with COVID-19 vaccine.

10. Have you received monoclonal antibodies or convalescent plasma for prevention or treatment of COVID-19 infection?

- » Vaccination with a dose of COVID-19 vaccine should be delayed for at least 90 days after treatment.

11. Do you have a bleeding disorder or are you taking any medications that could affect blood clotting?

- » If yes, apply direct pressure (without rubbing) to the injection site for 5 minutes or longer following immunization.

12. Have you ever felt faint or fainted after a past vaccination or medical procedure?

- » If yes, consider having them lie down for the immunization.

13. Did you have any side effects after a previous dose of COVID-19 vaccine?

- » If client had a serious or allergic reaction to a previous dose of COVID-19 vaccine, they should not be vaccinated until it is determined to be safe by an allergist or other health care provider.

14. Have you had a serious reaction to any other vaccine in the past?

- » If client had a serious or allergic reaction to any other vaccine in the past, they should not be vaccinated until it is determined to be safe by a Medical Health Officer.