

For the Period July 15 to 21, 2014

Haying continues in the province and livestock producers now have 42 per cent of the hay crop baled or put into silage, with an additional 49 per cent cut and ready for baling, according to Saskatchewan Agriculture's weekly Crop Report.

Rain showers and high humidity have delayed haying and decreased hay quality in some areas. Thirteen per cent is rated as excellent in quality, 77 per cent good, eight per cent fair and two per cent poor. Hay yields are slightly below the five-year average (2009-2013). The estimated average hay yields on dry land are reported as 1.3 tons per acre for alfalfa, 1.4 tons per acre for alfalfa/brome hay, 1.1 tons per acre for other tame hay, 0.9 tons per acre for wild hay and 1.7 tons per acre for greenfeed. On irrigated land, the estimated average hay yields are 1.9 tons per acre for alfalfa, 1.8 tons per acre for alfalfa/brome hay, 1.4 tons per acre for other tame hay and 2.3 tons per acre for wild hay and greenfeed.

Rain during the week ranged from trace amounts to 84 mm in the Nipawin area. Across the province, topsoil moisture on cropland is rated as 14 per cent surplus, 75 per cent adequate, 10 per cent short and one per cent very short. Hay land and pasture topsoil moisture is rated as 10 per cent surplus, 74 per cent adequate, 13 per cent short and three per cent very short. Some areas in the south are drier than normal and will soon need moisture to help crops mature and fill.

Warm weather has helped advance many crops and the majority are in fair to excellent condition.

Storms moved through the province last week, bringing strong winds, heavy rain and hail. Other sources of crop damage include insects such as grasshoppers and wheat midge and diseases such as root rot and leaf spots.

Farmers are busy haying and controlling insects and crop disease.

One year ago

Livestock producers had 69 per cent of the hay crop cut and 43 per cent baled or put into silage. Crops were in mostly good condition.

Follow the 2014 Crop Report on Twitter @SKAgriculture

For further information, contact Shannon Friesen, PAg,
Cropping Management Specialist, Moose Jaw, Regional Services Branch,
Toll Free: 1-866-457-2377 or 306-694-3592, E-mail: cropreport@gov.sk.ca.
Also available on the Ministry of Agriculture website at www.agriculture.gov.sk.ca.



Southeastern Saskatchewan (Crop District 1 – Carnduff, Estevan, Redvers, Moosomin and Kipling areas; Crop District 2 – Weyburn, Milestone, Moose Jaw, Regina and Qu’Appelle areas; Crop District 3ASE – Radville and Lake Alma areas)

Warm weather is helping to advance many crops in the region; however, many areas are in need of rain to help crops mature and fill. Rainfall varied across the region, ranging from trace amounts to 37 mm in the Moose Jaw area. The Moosomin area has received 600 mm of rain since April 1, the greatest amount for both the region and the province.

Estimated Southeast Hay Yields (tons/acre)		
July 21, 2014		
	Dry land	Irrigated Land
Alfalfa	1.3	N/A
Brome/Alfalfa	1.5	N/A
Other Tame Hay	1.2	N/A
Wild Hay	1.0	N/A
Greenfeed	1.7	N/A

Topsoil moisture conditions on cropland are rated as 15 per cent surplus, 72 per cent adequate and 13 per cent short. Hay land and pasture topsoil moisture is rated as 13 per cent surplus, 71 per cent adequate and 16 per cent short. Moisture conditions vary across the region. CD 1B is reporting that 29 per cent of the cropland has surplus topsoil moisture, while CDs 2A and 3ASE are reporting that 50 per cent of the cropland there is short topsoil moisture.

Livestock producers now have 37 per cent of the hay crop baled or put into silage, with an additional 23 per cent cut and ready for baling. Hay quality is rated as 11 per cent excellent, 82 per cent good, four per cent fair and three per cent poor.

The majority of crop damage is due to localized flooding, wind, hail and drought conditions. Producers are spraying for insects such as wheat midge and grasshoppers, and for diseases such as fusarium head blight, sclerotinia and leaf spots.

Farmers are busy haying, controlling pests and hauling grain.

Southwestern Saskatchewan (Crop District 3ASW – Coronach, Assiniboia and Ogema areas; Crop District 3AN – Gravelbourg, Mossbank, Mortlach and Central Butte areas; Crop District 3B – Kyle, Swift Current, Shaunavon and Ponteix areas; Crop District 4 – Consul, Maple Creek and Leader areas)

Although recent warm weather has helped crops to advance in the southwest, many of them are suffering from a lack of moisture and will soon need rain to mature and fill. Rainfall this week ranged from trace amounts to 37 mm in the Cabri area. Since April 1, the Cadillac area has received 385 mm of rain.

Topsoil moisture conditions are deteriorating in many areas of the region, particularly the Maple Creek area as well as areas near the Alberta and United States borders. Topsoil moisture conditions on cropland are rated as 70 per cent adequate, 23 per cent short and seven per cent very short. Hay land and pasture topsoil moisture is rated as 60 per cent adequate, 28 per cent short and 12 per cent short. CD 4A is reporting that 40 per cent of both cropland and hay land and pasture are very short topsoil moisture, while CD 4B is reporting that 20 per cent of cropland and 23 per cent of hay land and pasture is very short of topsoil moisture.

Haying continues and livestock producers now have 61 per cent baled or put into silage, with an additional 18 per cent cut and ready for baling. Hay quality is rated as 10 per cent excellent, 77 per cent good, 10 per cent fair and three per cent poor.

Estimated Southwest Hay Yields (tons/acre) July 21, 2014		
	Dry land	Irrigated Land
Alfalfa	1.2	1.8
Brome/Alfalfa	1.1	1.4
Other Tame Hay	0.8	N/A
Wild Hay	0.7	N/A
Greenfeed	1.2	2.6

Most crop damage this week is attributed to drought conditions. Other sources include localized flooding, hail, wind, insects and diseases. Some producers are spraying for grasshoppers and diseases in later-seeded crops.

Farmers are busy haying, controlling pests, fixing fences and hauling grain.

East-Central Saskatchewan (Crop District 5 – Melville, Yorkton, Cupar, Kamsack, Foam Lake, Preeceville and Kelvington areas; Crop District 6A – Lumsden, Craik, Watrous and Clavet areas)

Crops continue to recover from recent flooding, although in some areas many fields have been significantly affected. Rain will be needed in many areas of the region to help remaining crops mature and fill. Rainfall this past week ranged from trace amounts to 30 mm in the Saskatoon area. The Foam Lake area has received the most rainfall since April 1 (506 mm).

Topsoil moisture conditions on cropland are rated as 15 per cent surplus, 82 per cent adequate and three per cent short. Hay land and pasture topsoil moisture is rated as nine per cent surplus, 86 per cent adequate and five per cent short. CD 5B is reporting that 22 per cent of cropland and 14 per cent of hay land and pasture have surplus topsoil moisture.

Estimated East-central Hay Yields (tons/acre) July 21, 2014		
	Dry land	Irrigated Land
Alfalfa	1.5	2.1
Brome/Alfalfa	1.4	2.1
Other Tame Hay	1.3	1
Wild Hay	1.1	N/A
Greenfeed	1.8	1.5

Livestock producers now have 27 per cent of the hay crop baled or put into silage, with an additional 26 per cent cut and ready for baling. At this time, hay quality is rated as 19 per cent excellent, 75 per cent good and six per cent fair.

Hail, wind and localized flooding have caused the most damage this week. Insects such as wheat midge and grasshoppers are also causing some damage. Producers are spraying for sclerotinia, fusarium and leaf spot diseases.

Farmers are busy haying and controlling diseases and insects.

West-Central Saskatchewan (Crop Districts 6B – Hanley, Outlook, Loreburn, Saskatoon and Arelee areas; Crop District 7A – Rosetown, Kindersley, Eston, Major; CD 7B - Kerrobert, Macklin, Wilkie and Biggar areas)

Storms moved through the region late last week, bringing high winds, baseball-sized hail and heavy rain. Many fields suffered extensive hail damage, lodging and localized flooding. There are also reports of destroyed buildings, damaged vehicles and toppled grain bins.

Some areas received large amounts of rain within a couple of hours: the Biggar area received 82 mm. Since April 1, the Rosetown area has received 313 mm of rain, the most in the region.

Estimated West-central Hay Yields (tons/acre) July 21, 2014		
	Dry land	Irrigated Land
Alfalfa	1.4	2
Brome/Alfalfa	1.3	2.5
Other Tame Hay	1.0	2
Wild Hay	0.9	2.3
Greenfeed	1.7	3.2

Topsoil moisture conditions on cropland are rated as 15 per cent surplus, 79 per cent adequate and six per cent short. Hay land and pasture topsoil moisture is rated as 10 per cent surplus, 81 per cent adequate and nine per cent short. CD 6B is reporting that 12 per cent of cropland and 14 per cent of hay land and pasture is short of topsoil moisture. Rain would be welcomed in many areas of the region to help crops mature and fill.

Haying is progressing and livestock producers now have 41 per cent of the hay crop baled or put into silage, with an additional 30 per cent cut and ready for baling. Hay quality is rated as nine per cent excellent, 83 per cent good and eight per cent fair.

The majority of crop damage this week is attributed to hail, wind and localized flooding. There are also reports of damage from grasshoppers and from diseases such as leaf spot and root rot.

Farmers are busy haying, hauling grain and controlling pests.

Northeastern Saskatchewan (Crop District 8 – Hudson Bay, Tisdale, Melfort, Carrot River, Humboldt, Kinistino, Cudworth and Aberdeen areas; Crop District 9AE – Prince Albert, Choiceland and Paddockwood areas)

Some areas of the region received heavy rain last week that has flooded some fields and damaged crops. The Nipawin area reports receiving 84 mm of rain in one day, although there are reports of larger amounts near the Garrick area. Since April 1, the Arborfield area has received 345 mm of rain, the greatest amount for the region.

Estimated Northeast Hay Yields (tons/acre) July 21, 2014		
	Dry land	Irrigated Land
Alfalfa	1.5	N/A
Brome/Alfalfa	1.6	N/A
Other Tame Hay	1.0	N/A
Wild Hay	1.0	N/A
Greenfeed	1.0	N/A

Topsoil moisture conditions have worsened since last week and are now rated as 40 per cent surplus and 60 per cent adequate on cropland. Hay land and pasture topsoil moisture is rated as 67 per cent surplus and 33 per cent adequate. Many areas are in need of some warm and dry weather to dry up fields, while other areas will soon need a rain to help crops mature and fill.

Haying continues for livestock producers in the region as 35 per cent of the hay crop has been baled or put into silage. An additional 35 per cent is cut and ready for baling. Hay quality at this time is rated as eight per cent excellent, 67 per cent good, 17 per cent fair and eight per cent poor.

Crops are advancing quickly in the region, although wind and localized flooding have caused some damage. Producers are spraying for grasshoppers, wheat midge and diseases such as sclerotinia and leaf spots in later-seeded crops.

Farmers are busy haying, fixing fences and controlling pests.

Northwestern Saskatchewan (Crop District 9AW – Shellbrook, North Battleford, Big River and Hafford areas; Crop District 9B – Meadow Lake, Turtleford, Pierceland, Maidstone and Lloydminster areas)

Recent rain has helped crops advance, but it is slowing down haying. The Neilburg area is reporting that 38 mm of rain was received this past week, while the Hafford area is reporting 377 mm of rain since April 1, the greatest amount in the region.

Cropland topsoil moisture conditions are rated as three per cent surplus, 93 per cent adequate and four per cent short, while hay land and pasture topsoil moisture conditions are rated as one per cent surplus, 95 per cent adequate and four per cent short. For some in the region, rainfall would be welcomed, while for others, warm and dry weather is needed to help crops advance.

Thirty-six per cent of the hay crop has now been baled or put into silage, with an additional 26 per cent cut and ready for baling. At this time, hay quality is rated as 23 per cent excellent, 71 per cent good and six per cent fair.

Producers have been monitoring for insect pests and spraying for leaf disease and sclerotinia in later-seeded crops when weather allows. The majority of crop damage this past week was due to localized flooding, lodging and drought conditions in some areas.

Estimated Northwest Hay Yields (tons/acre) July 21, 2014		
	Dry land	Irrigated Land
Alfalfa	1.4	N/A
Brome/Alfalfa	1.5	N/A
Other Tame Hay	1.1	N/A
Wild Hay	1.0	N/A
Greenfeed	2.0	N/A


Farmers are busy haying, hauling grain and controlling pests.

SK (provincial) Crop Conditions - July 21, 2014							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	5	5	15	13	12	16	20
% good	59	49	63	67	59	59	45
% fair	25	43	18	16	25	19	17
% poor	10	2	3	4	3	5	12
% very poor	1	1	1	0	1	1	6
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	7	12	5	8	22	15	9
% good	57	56	78	66	58	55	81
% fair	27	23	16	21	15	21	10
% poor	6	8	1	5	4	4	0
% very poor	3	1	0	0	1	5	0
Southeast							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	4	20	12	14	8	14	22
% good	51	61	63	56	60	58	26
% fair	28	18	18	17	25	20	14
% poor	16	1	6	12	5	7	22
% very poor	1	0	1	1	2	1	16

	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	7	10	1	9	7	2	0
% good	47	59	71	64	45	32	60
% fair	29	22	25	22	27	33	40
% poor	10	8	2	5	12	13	0
% very poor	7	1	1	0	9	20	0
Southwest							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	0	0	8	12	3	16	3
% good	88	37	70	72	72	65	80
% fair	10	62	21	15	25	18	15
% poor	2	1	1	1	0	1	2
% very poor	0	0	0	0	0	0	0
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	0	6	1	0	23	15	8
% good	78	80	85	60	57	65	82
% fair	22	13	13	40	15	18	10
% poor	0	1	1	0	3	2	0
% very poor	0	0	0	0	2	0	0
East-central							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	9	6	9	15	5	7	8
% good	47	42	59	60	56	57	58
% fair	32	37	26	21	31	28	22
% poor	9	11	5	4	6	7	12
% very poor	3	4	1	0	2	1	0
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	9	5	3	0	16	14	0
% good	53	45	54	75	47	53	75
% fair	31	37	32	13	28	21	25
% poor	6	12	11	12	8	10	0
% very poor	1	1	0	0	1	2	0
West-central							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	16	7	24	16	22	24	35


% good	74	73	68	68	65	63	50
% fair	10	20	8	16	13	12	15
% poor	0	0	0	0	0	1	0
% very poor	0	0	0	0	0	0	0
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	24	22	20	5	21	27	60
% good	68	66	63	75	71	56	30
% fair	8	12	17	20	8	16	10
% poor	0	0	0	0	0	1	0
% very poor	0	0	0	0	0	0	0
Northeast							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	0	0	11	0	11	10	0
% good	100	100	58	55	55	55	59
% fair	0	0	27	35	31	23	23
% poor	0	0	3	8	3	11	17
% very poor	0	0	1	2	0	1	1
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	7	9	0	5	1	0	N/A
% good	57	51	25	50	52	70	N/A
% fair	33	24	65	30	20	18	N/A
% poor	3	14	5	10	21	7	N/A
% very poor	0	2	5	5	6	5	N/A
Northwest							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	28	1	33	N/A	31	30	N/A
% good	37	99	57	N/A	62	63	N/A
% fair	35	0	10	N/A	7	7	N/A
% poor	0	0	0	N/A	0	0	N/A
% very poor	0	0	0	N/A	0	0	N/A
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	30	28	N/A	N/A	31	0	N/A
% good	70	56	N/A	N/A	51	75	N/A
% fair	0	15	N/A	N/A	17	25	N/A
% poor	0	1	N/A	N/A	1	0	N/A
% very poor	0	0	N/A	N/A	0	0	N/A

Crop Districts and Rural Municipalities in Saskatchewan



Weekly Rainfall

for the week ending July 21, 2014



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Weekly Rainfall Summary

(in millimeters)

1 inch = 25 mm

for the period July 15 to 21, 2014


Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr	Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr	Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr	
1A	2	Mount Pleasant	5	233	4A	49	White Valley	5	153	7A	259	Snipe Lake	N/A	20	
	3	Enniskillen	5	186		51	Reno	18.4	115.7		287	St. Andrews	31	313	
	33	Moose Creek	N/A	244		79	Arlington	9	264		288	Pleasant Valley	27	257	
	34	Browning	N/A	240		109 A	Carmichael	N/A	181		290 A	Kindersley	17	167	
	61	Antler	N/A	432		109 B	Carmichael	N/A	149		290 B	Kindersley	N/A	105.5	
	63	Moose Mountain	3	345		110	Piapot	NIL	135		290 C	Kindersley	22	159.2	
	64	Brook	3	245		111	Maple Creek	NIL	93		292	Milton	29	218	
	65	Tecumseh	5	329		4B	139	Gull Lake	6		207	317 A	Marriott	N/A	127.5
	91	Maryfield	4	457			141	Big Stick	NIL		140	317 B	Marriott	81	278
	122	Martin	NIL	600			142	Enterprise	NIL		184	318	Mountain View	56	297
123	Silverwood	NIL	332	169	Pittville		N/A	162	320 A	Oakdale	29	197.1			
124	Kingsley	NIL	320	231	Happyland		27	175	320 B	Oakdale	45	199			
125 A	Chester	NIL	318	5A	183 A		Fertile Belt	NIL	344	321	Prairiedale	23	220		
125 B	Chester	NIL	342		183 B		Fertile Belt	NIL	369	347	Biggar	40	292		
151 A	Rocanville	NIL	424		186		Abernethy	2	360	350 A	Mariposa	82	211.6		
151 B	Rocanville	NIL	503		211		Churchbridge	NIL	305	350 B	Mariposa	77	295		
154	Elcapo	NIL	375		213		Saltcoats	1	383	351	Progress	75	240		
155 A	Wolseley	0.5	304.5		216	Tullymet	N/A	272.5	352	Heart's Hill	49	204			
155 B	Wolseley	N/A	378		241	Calder	4	269	377	Glenside	N/A	289			
67	Weyburn	2	198		243	Wallace	NIL	319	378 A	Rosemount	55	245			
68	Brokenshell	NIL	200		244	Orkney	N/A	415	378 B	Rosemount	34	285			
97	Wellington	NIL	214.5		245 A	Garry	2	400	379	Reford	53	311			
2B	127 A	Francis	2	316	245 B	Garry	NIL	435	381	Grass Lake	N/A	206			
	127 B	Francis	0.3	225.7	245 C	Garry	2	432	382	Eye Hill	62.5	231.6			
2B	129	Bratt's Lake	4	239.5	246	Ituna Bon Accord	2	371	409	Buffalo	30	221			
	131 A	Baildon	1	284	247	Kelross	3	335	410	Round Valley	40	285.4			
	131 B	Baildon	10	362.2	248	Touchwood	2	295	8A	395	Porcupine	13	243		
	156 A	Indian Head	NIL	372.5	271	Cote	1	277		397	Barrier Valley	13.1	304.3		
	156 B	Indian Head	NIL	408.5	273	Sliding Hills	N/A	477		428	Star City	16	340		
	157	South Qu'Appelle	NIL	312	277	Emerald	5	506		456	Arborfield	40	345		
	160 A	Pense	NIL	299	305	Invermay	NIL	400		457	Connaught	33	265		
	160 B	Pense	N/A	75	307	Elfros	10	389		486	Moose Range	84	285		
	161	Moose Jaw	4	372	308 A	Big Quill	10	292		487	Nipawin	N/A	214		
	162	Caron	37	363	308 B	Big Quill	3	268		8B	369	St. Peter	21	300	
191	Marquis	NIL	318	331	Livingston	N/A	306	370 A			Humboldt	28	331		
38 A	Laurier	7	259.9	334	Preeceville	N/A	112	370 B			Humboldt	N/A	306		
38 B	Laurier	9	244	336	Sasman	3	293	371	Bayne		32	343			
39 A	The Gap	10	190	337	Lakeview	6	294	372	Grant		22	328.5			
39 B	The Gap	6	216	338	Lakeside	13	313	400	Three Lakes		12	342			
3ASW	10	Happy Valley	2	156	366	Kelvington	10	327	402		Fish Creek	20	315		
	12	Poplar Valley	N/A	256	367	Ponass Lake	25	314	429		Flett's Springs	12	268		
	40 A	Bengough	3	191	6A	190 A	Dufferin	3	480		459	Kinistino	10	308	
	40 B	Bengough	NIL	224		190 B	Dufferin	9	386.5		460	Birch Hills	13.7	282.9	
	42	Willow Bunch	2	232		190 C	Dufferin	NIL	335	9AE	488	Torch River	50	266	
	43	Old Post	3	260		190 D	Dufferin	1	189.5		491	Buckland	N/A	317	
	70	Key West	NIL	63.6		219	Longlaketon	1	301		9AW	406	Mayfield	34	263
	73 A	Stonehenge	3.8	246.3		220	Mckillop	2	435			435	Redberry	30	377
	73 B	Stonehenge	8	316		221	Sarnia	2.9	381.7			436	Douglas	30	313
	74	Wood River	NIL	175.5		222	Craik	NIL	320	463		Duck Lake	35	314.9	
101	Terrell	5	248	251		Big Arm	6	414	9B	467 A		Round Hill	30	362	
102	Lake Johnston	20.3	215.4	252		Arm River	1	370		467 B	Round Hill	27	205		
103	Sutton	NIL	150	279	Mount Hope	4	330	438		Battle River	30	187			
132 A	Hillsborough	5	223	282	McCraney	5	362	440		Hillsdale	38	294.5			
132 B	Hillsborough	6	291	309	Prairie Rose	N/A	350	442		Manitou Lake	34.3	216.1			
134	Shamrock	N/A	222	310	Usborne	12	166.5	498 A		Parkdale	26	243.5			
193 A	Eyebrow	1	301	312	Morris	10	113	498 B		Parkdale	27	200			
193 B	Eyebrow	N/A	295	313	Lost River	N/A	306	499		Mervin	35	191.5			
224	Maple Bush	N/A	294	339	Leroy	13	380.1	501 A		Frenchman Butte	27	268			
3BS	17	Val Marie	8	385	340	Wolverine	26	323		501 B	Frenchman Butte	30	214		
	75 A	Pinto Creek	NIL	248	341	Viscount	15	287	501 C	Frenchman Butte	31	259			
	75 B	Pinto Creek	NIL	220	343 A	Blucher	20	282.9	502	Britannia	27.5	184.5			
	76	Auvergne	N/A	185	343 B	Blucher	30	170	561	Loon Lake	35	277			
	77	Wise Creek	5	288	254	Loreburn	N/A	246	588 A	Meadow Lake	25	272			
	78	Grassy Creek	18	270	284	Rudy	17	298	588 B	Meadow Lake	20	303			
	105	Glenbain	1	179	285	Fertile Valley	27	224	588 C	Meadow Lake	29	255			
	106	Whiska Creek	N/A	303	286	Milden	N/A	153	588 D	Meadow Lake	13.8	321.2			
	107	Lac Pelletier	N/A	170	314	Dundurn	27	266	622	Beaver River	N/A	191.2			
	108	Bone Creek	N/A	206	344	Corman Park	32	295							
137	Swift Current	N/A	41.6	346	Perdue	31	283								
38A	138 A	Webb	17	221	376	Eagle Creek	27	292							
	138 B	Webb	17.5	230.3	403	Rosthern	34	254							
	166	Excelsior	9	207											
	167	Sask. Landing	3.4	214.4											
	168 A	Riverside	6	192											
	168 B	Riverside	37.1	201.2											
	226	Victory	10	227											
	228	Lacadena	15.5	185											
	257	Monet	13	138.6											

These precipitation amounts represent point locations within each municipality and do not necessarily reflect the whole R. M.

Cumulative Rainfall

From: April 1, 2014


To: July 21, 2014



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Cropland Topsoil Moisture Conditions


July 22, 2014



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Hay and Pasture Topsoil Moisture Conditions

July 22, 2014



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.