

## For the Period July 19 to 25, 2016

Good progress was made with haying this past week. Livestock producers have 50 per cent of the hay crop baled or put into silage and another 20 per cent cut and ready for baling, according to Saskatchewan Agriculture's weekly Crop Report. The five-year average (2011-2015) is 55 per cent baled or put into silage.

Hay quality is rated as five per cent excellent, 70 per cent good, 24 per cent fair and one per cent poor. Hay is slow to dry in the swath due to rain and high humidity.

Crops are generally in good condition. Sixty-two per cent of the spring wheat, winter wheat, flax and canola are in good condition. Wet conditions across much of the lentil-producing area has affected quality. Lentils are rated as 15 per cent excellent, 47 per cent good, 26 per cent fair, 10 per cent poor and two per cent very poor. Peas are rated as 30 per cent excellent, 52 per cent good, 16 per cent fair and two per cent poor.

Most areas of the province received rain during the past week, with the heaviest showers dropping between 50 and 100 mm of precipitation. There were reports of 200 mm of rain in the Elfros region. Heavy rain over the past several weeks has caused crops to lodge, and they remain under flooding stress in some areas. Lentils and peas in many areas of the province are suffering from too much moisture. Diseases and hail have also caused crop damage.

Provincially, cropland topsoil moisture is rated as 15 per cent surplus, 80 per cent adequate and five per cent short. Hay land and pasture topsoil moisture is rated as nine per cent surplus, 83 per cent adequate and eight per cent short.

Harvesting of winter wheat, peas and lentils in some areas is expected to start in the coming week. Farmers are busy haying and getting ready for harvest.

### One year ago

Sixty-seven of the hay crop had been baled or put into silage. Harvest was just beginning with desiccation of pulses and some winter cereals combined. Moisture conditions improved with recent rains.

Follow the 2016 Crop Report on Twitter at @SKAgriculture

For further information, contact Daphne Cruise, PAg,  
Crop Management Specialist, Moose Jaw, Regional Services Branch,  
Toll Free: 1-866-457-2377 or 306-694-3587, E-mail: [cropreport@gov.sk.ca](mailto:cropreport@gov.sk.ca).  
Also available on the Ministry of Agriculture website at [www.saskatchewan.ca/crop-report](http://www.saskatchewan.ca/crop-report).

### **Southeastern Saskatchewan:**

- Crop District 1 – Carnduff, Estevan, Redvers, Moosomin and Kipling areas
- Crop District 2 – Weyburn, Milestone, Moose Jaw, Regina and Qu'Appelle areas
- Crop District 3ASE – Radville and Lake Alma areas

Fifty per cent of the hay crop is baled and 22 per cent is cut and ready for baling. Haying is furthest advanced in Crop District 2A, which has 65 per cent baled, and least advanced in Crop District 1B, which has 36 per cent baled. Hay quality is rated as 79 per cent good and 21 per cent fair. High humidity and rain showers are causing quality loss in the swath. Dry weather is needed to get the rest of the hay up in good shape. There are a few reports of producers starting to take their second cut of hay.

The southeast recorded varying amounts of rain during the week, although amounts were less than they have been in the previous weeks. Most areas, with the exception of the few that were hit with thunderstorms, reported less than 15 mm of rain. The Carnduff area reported 15 mm, the Grenfell area 28 mm, Weyburn 34mm, Moose Jaw 73 mm and Radville 10 mm. The Maryfield area holds the regional record for greatest amount of precipitation since April 1 (413 mm). The Odessa area has recorded the least amount of rain for the region since April 1 (210 mm).

Cropland topsoil moisture is rated as 16 per cent surplus, 81 per cent adequate and three per cent short. Hay land and pasture topsoil moisture is rated as 10 per cent surplus, 85 per cent adequate and five per cent short. Crop districts 1B and 2B are reporting that 32 and 23 per cent, respectively, of cropland acres have surplus topsoil moisture at this time. Twenty-nine per cent of hay land and pasture in CD 1B has surplus topsoil moisture. There is crop loss where land has flooded or remained wet.

Most crops in the region remain in good condition, although the pulse crops are succumbing to excess moisture stress. Crop damage this week was mainly caused by localized flooding and diseases such as root rot, leaf spots, sclerotinia and botrytis. Hail was reported in the region, which caused significant damage in some areas. Marble- to golf-ball-sized hail was reported in the Indian Head and Moose Jaw regions. Some producers are expecting to start harvest in the next week if the weather cooperates.

### **Southwestern Saskatchewan:**

- Crop District 3ASW – Coronach, Assiniboia and Ogema areas
- Crop District 3AN – Gravelbourg, Mossbank, Mortlach and Central Butte areas
- Crop District 3B – Kyle, Swift Current, Shaunavon and Ponteix areas
- Crop District 4 – Consul, Maple Creek and Leader areas

Scattered and sporadic rain showers in the region allowed some producers to make good progress in haying while others were slowed. Fifty per cent of the hay crop has been baled and an additional 19 is cut and ready for baling. Hay quality is rated as three per cent excellent, 66 per cent good and 31 per cent fair. Haying is most advanced in Crop District 4A, which has 65 per cent baled, and least advanced in Crop District 3AW, which has 30 per cent baled. The hay has been slow to dry due to frequent rain showers and high humidity. Rainfall varied greatly in the region, from nothing to 93 mm in the Kyle area. Other areas receiving large amounts of rain include Moose Jaw (64 mm), Big Beaver (30

mm), Shaunavon (28 mm), Swift Current (31 mm) and Leader (52 mm). The Webb area holds the regional record for greatest amount of precipitation since April 1 (433 mm).

Cropland topsoil moisture conditions are rated as 14 per cent surplus, 84 per cent adequate and two per cent short. Hay land and pasture topsoil moisture is rated as six per cent surplus, 89 per cent adequate and five per cent short. Crop District 4B is reporting that 25 per cent of the cropland has surplus topsoil moisture. Crop loss is occurring due to excess moisture in some areas, particularly in low spots and areas in the field that may have been saturated prior to the rain.

The majority of crops in the region are in good condition, although a pulse quality downgrade is imminent due to excess moisture in areas. The majority of crop damage this past week is attributed to localized flooding, strong winds, lodging and hail. There are reports of some roads being washed out in the Kyle area. There are a couple reports of fall rye being combined. Although harvest will be later than anticipated because of the moisture, some pea crops are expected to be harvested in the coming week.

#### **East-Central Saskatchewan:**

- Crop District 5 – Melville, Yorkton, Cupar, Kamsack, Foam Lake, Preeceville and Kelvington areas
- Crop District 6A – Lumsden, Craik, Watrous and Clavet areas

Good haying progress was made in some areas of the region. Forty-five per cent of the hay crop has been baled and an additional 23 per cent is cut and ready to bale. Crop District 6A has 53 per cent baled, while Crop District 5B has 38 per cent baled. Hay quality is rated as seven per cent excellent, 74 per cent good, 15 per cent fair and four per cent poor. A few days of warm and breezy conditions allowed haying to continue toward the end of the week. Haying has been very slow in the region due to less-than-ideal weather conditions. Frequent rain showers and high humidity have caused quality to deteriorate.

Heavy rain fell on the region during the past week. Amounts ranged from 4 mm (Bulyea area) to 93 mm (Wynyard area). There were reports of over 200 mm falling on the Elfron region within two hours. Roads and fields were flooded and culverts were washed out. Most other areas recorded over 25 mm of rain for the week. The Kelvington area holds the regional record for greatest amount of precipitation since April 1 (365 mm).

Topsoil moisture conditions on cropland are rated as 17 per cent surplus and 83 per cent adequate. Hay land and pasture topsoil moisture is rated as 13 per cent surplus, 86 per cent adequate and one per cent short. Crop District 5B is reporting that 30 per cent of the cropland has surplus topsoil moisture at this time. There is crop loss due to excess moisture in some areas, particularly in low spots and areas in the field that have been saturated for a couple of weeks.

Most crops remain in good condition. Sources of crop damage this week include localized flooding, hail, wind and diseases such as root rot and other pulse diseases. Some pulse crops are expected to be desiccated in the coming week if weather cooperates.

### **West-Central Saskatchewan:**

- Crop Districts 6B – Hanley, Outlook, Loreburn, Saskatoon and Arelee areas
- Crop District 7A – Rosetown, Kindersley, Eston, Major
- Crop District 7B - Kerrobert, Macklin, Wilkie and Biggar areas

Haying continues despite the rain. Forty-seven per cent of the hay crop has been baled or put into silage, and 24 per cent is cut and ready for baling. Continuous rain and high humidity are taking their toll on hay quality, which is now rated as five per cent excellent, 48 per cent good and 47 per cent fair.

Rainfall varied greatly throughout the region, ranging from nothing in the Tramping Lake and Unity areas to 50 mm in the Conquest area. Other areas that received large amounts of rain this past week include Dinsmore (37 mm), Sonningdale (24 mm), Kindersley (48 mm), Biggar (41 mm) and Macklin (37 mm). The Conquest area holds the regional record for greatest amount of precipitation since April 1 (332 mm).

Topsoil moisture conditions on cropland are rated as 15 per cent surplus and 83 per cent adequate and two per cent short. Hay land and pasture topsoil moisture is rated as 10 per cent surplus, 86 per cent adequate and four per cent short. Crop District 7A is reporting that 25 per cent of cropland has surplus topsoil moisture. There is crop loss due to excess moisture in some areas.

The majority of crops are in good condition, but pulses are taking the biggest hit when it comes yield loss due to excess moisture. Crop damage this week is attributed to localized flooding, diseases and hail. Baseball-sized hail and heavy rain was reported in the Outlook and Conquest areas, causing significant damage to crops and property. There are predictions of peas being desiccated this coming week if weather cooperates. Warm and dry weather is needed as harvest approaches if fields in some areas are to be accessed. Despite the excess moisture in most areas of the region, there are few areas that could use more moisture to help fill and finish off the crops.

### **Northeastern Saskatchewan:**

- Crop District 8 – Hudson Bay, Tisdale, Melfort, Carrot River, Humboldt, Kinistino, Cudworth and Aberdeen areas
- Crop District 9AE – Prince Albert, Choiceland and Paddockwood areas

The region is furthest ahead in haying, with 59 per cent of the hay crop baled or put into silage. An additional 17 per cent is cut and ready for baling. Ninety-two per cent of the hay is rated as good quality and eight per cent is rated as poor. Although rain was reported, it resulted in very little down time for cutters and balers in most areas. High humidity is causing delays in baling.

Rainfall ranged from nothing in the Kinistino area to 36 mm in the Porcupine Plain and Tisdale areas. Other areas receiving significant rainfall include Star City (20 mm), Birch Hills (15 mm), Alvena (6 mm), and Garrick (21 mm). The Porcupine Plain area holds the regional record for greatest amount of precipitation since April 1 (348 mm).

Cropland topsoil moisture conditions are rated as 27 per cent surplus, 71 per cent adequate and two per cent short. Hay land and pasture topsoil moisture is rated as 36 per

cent surplus and 64 per cent adequate. Crop District 8A is reporting that 46 per cent of cropland and 43 per cent of hay land and pasture have surplus topsoil moisture at this time.

Crops remain are generally in good condition. Crop damage includes hail, lodging and disease. Significant rain in the past few weeks has washed out roads in some areas and is flooding crops, causing loss. Some harvesting will begin in the next week or so. Warm, dry weather is needed to dry the soil enough to support harvest equipment.

#### **Northwestern Saskatchewan:**

- Crop District 9AW – Shellbrook, North Battleford, Big River and Hafford areas
- Crop District 9B – Meadow Lake, Turtleford, Pierceland, Maidstone and Lloydminster areas

Drier weather has allowed for good progress to be made in the hay field. Fifty-three per cent of the hay crop has been baled and another 21 per cent is cut and ready for baling. Hay quality at this time is rated as 14 per cent excellent, 72 per cent good and 14 per cent fair. The region did not receive as much rain as other areas in the province, so haying was able to continue for the most part. Hay that has been cut and baled the last two weeks is of much better quality as little, if any, rain fell on the swaths.

Rainfall ranged from nil to 16 mm (Turtleford area). The Hafford and Lloydminster areas received 12 mm, Debden 7 mm and Barthel and Meadow Lake 10 mm. The Frenchman Butte area holds the regional record for greatest amount of precipitation since April 1 (278 mm). The Speers area has reported the least amount of rainfall since April 1 (100 mm).

Cropland topsoil moisture is rated as one per cent surplus, 71 per cent adequate and 28 per cent short. Hay land and pasture topsoil moisture is rated as 66 per cent adequate, 33 per cent short and one per cent very short. Although most of the region has adequate topsoil moisture, there are areas that are short on moisture and could use a good rain.

Crops in general are in good condition. The region has not received as much precipitation throughout the growing season as the rest of the province. Crop damage this past week is attributed to strong winds, lodging and diseases, particularly in peas.

<b>Provincial</b>							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	36	48	31	37	23	27	19
% good	61	50	62	58	69	65	73
% fair	3	2	7	4	7	7	6
% poor	0	0	0	1	1	1	1
% very poor	0	0	0	0	0	0	1
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	31	31	27	9	30	15	8
% good	62	61	66	66	52	47	68
% fair	6	6	6	20	16	26	8
% poor	1	1	1	5	2	10	16
% very poor	0	1	0	0	0	2	0

<b>Southeast</b>							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	48	46	39	35	42	32	23
% good	51	50	56	62	50	54	71
% fair	1	4	5	3	7	13	5
% poor	0	0	0	0	1	1	1
% very poor	0	0	0	0	0	0	0
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	35	36	10	8	35	17	N/A
% good	60	59	90	64	55	49	N/A
% fair	4	5	0	23	7	21	N/A
% poor	1	0	0	5	2	8	N/A
% very poor	0	0	0	0	1	5	N/A

<b>Southwest</b>							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	38	57	37	39	41	27	25
% good	57	43	58	56	57	70	61
% fair	5	0	5	5	2	3	10
% poor	0	0	0	0	0	0	3
% very poor	0	0	0	0	0	0	1
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	26	44	28	37	25	10	25
% good	68	51	62	63	48	47	61
% fair	5	4	8	0	26	29	10
% poor	1	1	2	0	1	13	3
% very poor	0	0	0	0	0	1	1

East-central							
	Winter whea	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	29	27	33	33	28	31	9
% good	69	70	59	56	66	62	71
% fair	2	3	7	11	6	6	20
% poor	0	0	1	0	0	1	0
% very poor	0	0	0	0	0	0	0

East-central (continued)							
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	36	33	N/A	10	35	34	N/A
% good	58	58	N/A	78	55	38	N/A
% fair	5	7	N/A	10	8	25	N/A
% poor	1	2	N/A	1	1	3	N/A
% very poor	0	0	N/A	1	1	0	N/A

West-central							
	Winter whea	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	20	N/A	36	36	35	33	25
% good	73	N/A	57	61	56	61	73
% fair	7	N/A	7	3	9	6	2
% poor	0	N/A	0	0	0	0	0
% very poor	0	N/A	0	0	0	0	0
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	23	38	37	20	34	15	N/A
% good	65	58	59	54	50	45	N/A
% fair	11	4	4	26	9	24	N/A
% poor	1	0	0	0	6	11	N/A
% very poor	0	0	0	0	1	5	N/A



Northeast							
	Winter whea	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	N/A	N/A	18	45	19	33	16
% good	N/A	N/A	73	53	74	61	66
% fair	N/A	N/A	9	2	7	6	12
% poor	N/A	N/A	0	0	0	0	6
% very poor	N/A	N/A	0	0	0	0	0
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	16	16	N/A	N/A	16	N/A	N/A
% good	66	71	N/A	N/A	66	N/A	N/A
% fair	17	7	N/A	N/A	14	N/A	N/A
% poor	1	4	N/A	N/A	2	N/A	N/A
% very poor	0	2	N/A	N/A	2	N/A	N/A

Northwest							
	Winter whea	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	N/A	N/A	22	N/A	7	11	N/A
% good	N/A	N/A	70	N/A	84	76	N/A
% fair	N/A	N/A	8	N/A	9	13	N/A
% poor	N/A	N/A	0	N/A	0	0	N/A
% very poor	N/A	N/A	0	N/A	0	0	N/A
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	N/A	25	N/A	N/A	32	N/A	N/A
% good	N/A	67	N/A	N/A	59	N/A	N/A
% fair	N/A	8	N/A	N/A	8	N/A	N/A
% poor	N/A	0	N/A	N/A	1	N/A	N/A
% very poor	N/A	0	N/A	N/A	0	N/A	N/A



# Crop Districts and Rural Municipalities in Saskatchewan

- Regional Service Office
- Regional Satellite Office
- ▭ Crop District
- ▭ Rural Municipality




Data Source:  
Crop Districts - Saskatchewan Ministry of Agriculture

Geomatic Services, Ministry of Agriculture June 10, 2014

# Weekly Rainfall


from July 19 to July 25, 2016



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

# Weekly Rainfall

from July 19 to July 25, 2016



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.



# Weekly Rainfall Summary

(in millimeters)

1 inch = 25 mm

for the period 19-25, 2016


Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr	Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr	Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr
1A	2	Mount Pleasant	15	294	4A	49	White Valley	7	272	7A	287	St. Andrews	31	308
	3	Enniskillen	3	273		51	Reno	NIL	164		288	Pleasant Valley	N/A	166
	33	Moose Creek	N/A	239		79	Arlington	11	326		290 A	Kindersley	25	198
	34	Browning	N/A	335		109 A	Carmichael	N/A	281		290 B	Kindersley	48	254
	61	Antler	N/A	282		109 B	Carmichael	N/A	216		290 C	Kindersley	35	239
	64	Brock	2	234		110	Piapot	3	282		292	Milton	31	314
1B	65	Tecumseh	11	236	111	Maple Creek	N/A	208	317 A	Marriott	N/A	220		
	91	Maryfield	6	413	4B	139	Gull Lake	NIL	320	317 B	Marriott	16	191	
	122	Martin	15	325		142	Enterprise	N/A	108	318	Mountain View	41	251	
	123	Silverwood	7	278		169	Pittville	N/A	282	320 A	Oakdale	10	219	
	124 A	Kingsley	9	286		231	Happyland	52	336	320 B	Oakdale	9	264	
	124 B	Kingsley	N/A	NIL	5A	183	Fertile Belt	N/A	311	321	Prairieedale	20	280	
	125 A	Chester	10	375		211	Churchbridge	N/A	223	7B	347	Biggar	25	180
	125 B	Chester	21	294		213	Saltcoats	33	162		350 A	Mariposa	NIL	147
	151	Rocanville	10	312		241	Calder	36	189		350 B	Mariposa	N/A	146
	154	Elcapo	N/A	280		243	Wallace	N/A	215		351	Progress	N/A	195
155	Wolseley	28	322	244		Orkney	N/A	186	352		Heart's Hill	18	326	
2A	67	Weyburn	7	252	245 A	Garry	46	248	377		Glenside	7	162	
	68	Brokenshell	9	313	245 B	Garry	25	196	378	Rosemount	10	187		
2B	97	Wellington	N/A	253	245 C	Garry	N/A	148	379	Reford	12	175		
	127 A	Francis	10	210	246	Ituna Bon Accord	12	155	381	Grass Lake	4	144		
	127 B	Francis	21	212	247	Kellross	7	161	382	Eye Hill	37.5	249		
	129	Bratt's Lake	10	247	248	Touchwood	71	229	409	Buffalo	N/A	125		
	131 A	Baildon	N/A	304	5B	271	Cote	N/A	170	410	Round Valley	NIL	136	
	131 B	Baildon	28	375		273	Sliding Hills	8	128	8A	395	Porcupine	36	348
	156 A	Indian Head	36.7	271		277	Emerald	32	232		397	Barrier Valley	19.8	284
	156 B	Indian Head	21	293		305	Invermay	59	319		428	Star City	20	275
	159	Sherwood	34	230		307	Elfros	58	324		456	Arborfield	18	346
	160	Pense	12	240		308 A	Big Quill	93	253		457	Connaught	36	265
	161	Moose Jaw	N/A	271	308 B	Big Quill	N/A	184	486		Moose Range	16	291	
	3ASE	162	Caron	20	284	331	Livingston	36	292	487	Nipawin	21	310	
		191	Marquis	19	276	336	Sasman	27	227	8B	369	St. Peter	9	166
		38 A	Laurier	10.4	356	337	Lakeview	N/A	264		370 A	Humboldt	N/A	205
38 B		Laurier	5	284	338	Lakeside	34	270	370 B	Humboldt	N/A	186		
3ASW	39	The Gap	N/A	291	366	Kelvington	39	365	371	Bayne	4	252		
	10	Happy Valley	30.5	315	367	Ponass Lake	10	273	372	Grant	3.4	131		
3AN	12	Poplar Valley	2	249	6A	190 A	Dufferin	26	321	400	Three Lakes	5	222	
	40	Bengough	N/A	N/A		190 B	Dufferin	15	289	402	Fish Creek	6	174	
	42	Willow Bunch	NIL	297		190 C	Dufferin	37	260	429	Flett's Springs	5	155	
	43	Old Post	NIL	259		190 D	Dufferin	N/A	192	459	Kinistino	NIL	164	
	73 A	Stonehenge	NIL	332		219 A	Longlaketon	N/A	196	460	Birch Hills	14.8	166	
	73 B	Stonehenge	N/A	348		219 B	Longlaketon	28	241	9AE	488	Torch River	21	272
	102	Lake Johnston	NIL	211		220	McKillop	4	219		520	Paddockwood	N/A	143
	103	Sutton	5	267		221 A	Sarnia	10.7	274	521	Lakeland	N/A	143	
3BS	132 A	Hillsborough	9	287	221 B	Sarnia	N/A	222	9AW	406	Mayfield	8	170	
	132 B	Hillsborough	8	319	222	Craik	19	288		435	Redberry	12	173	
	134	Shamrock	N/A	17	251	Big Arm	20	217	436	Douglas	NIL	100		
	193 A	Eyebrow	7	260	252	Arm River	N/A	209	463	Duck Lake	N/A	136		
	193 B	Eyebrow	N/A	340	279	Mount Hope	N/A	233	467 A	Round Hill	NIL	220		
	17	Val Marie	9.3	304	282	McCraney	59	275	467 B	Round Hill	2	186		
	18	Lone Tree	17	358	312	Morris	48	344	494	Canwood	7	124		
	75	Pinto Creek	4	354	313	Lost River	16	185	9B	438	Battle River	4	184	
	76	Auvergne	3	273	339	Leroy	31	207		440	Hillsdale	NIL	99	
	77	Wise Creek	4	350	340	Wolverine	23	244		442	Manitou Lake	NIL	187	
78	Grassy Creek	3.5	265	341	Viscount	20	196	498 A		Parkdale	3	194		
105	Glenbain	7	347	343 A	Blucher	N/A	149	498 B		Parkdale	1	41		
106	Whiska Creek	11	319	343 B	Blucher	12	157	499 A		Mervin	16.1	207		
3BN	107	Lac Pelletier	N/A	340	6B	223 A	Huron	N/A	236	499 B	Mervin	N/A	N/A	
	108	Bone Creek	28	353		223 B	Huron	5	262	501 A	Frenchman Butte	8	184	
	138 A	Webb	NIL	433		284 A	Rudy	46	286	501 B	Frenchman Butte	5	177	
	138 B	Webb	4	319		285 B	Rudy	32	284	501 C	Frenchman Butte	N/A	278	
	166	Excelsior	24	341		285 A	Fertile Valley	50	332	502	Britannia	12	237	
	167	Sask. Landing	8.2	330		285 B	Fertile Valley	46	323	561	Loon Lake	10	187	
	168 A	Riverside	31	369		286	Mildern	37	298	588 A	Meadow Lake	12	154	
	168 B	Riverside	17.8	234		314	Dundurn	N/A	206	588 B	Meadow Lake	10	164	
	226	Victory	14	276		344	Corman Park	N/A	111	622	Beaver River	12.4	184	
	228	Lacadena	24	329		346	Perdue	21	280					
257	Monet	93	433	376	Eagle Creek	24	196							
				403	Rosthern	2	140							

These precipitation amounts represent point locations within each municipality and do not necessarily reflect the whole R. M.

Municipality No: A, B, C and D - more than one reporter

# Cumulative Rainfall


from April 1 to July 25, 2016



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

# Cropland Topsoil Moisture Conditions

July 25, 2016




NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.




# Hay and Pasture Topsoil Moisture Conditions

## July 25, 2016



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.



Data Source:  
Moisture - Ministry of Agriculture, Crop Report Database  
IDW interpolation (power 2.5, fixed radius 300 km)

Geomatics Services, Ministry of Agriculture    July 27, 2016